

A POWER MEAN INEQUALITY INVOLVING THE COMPLETE ELLIPTIC INTEGRALS

GENDI WANG, XIAOHUI ZHANG, AND YUMING CHU

ABSTRACT. In this paper the authors investigate a power mean inequality for a special function which is defined by the complete elliptic integrals.

KEYWORDS: complete elliptic integrals, power mean, inequality
2010 MSC: 33E05, 30C62

1. INTRODUCTION

Throughout this paper we let $r' = \sqrt{1 - r^2}$ for $0 < r < 1$. The well-known complete elliptic integrals of the first and second kind [9, 11] are respectively defined by

$$(1.1) \quad \begin{cases} \mathcal{K}(r) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - r^2 \sin^2 \theta}}, \\ \mathcal{K}'(r) = \mathcal{K}(r'), \\ \mathcal{K}(0) = \frac{\pi}{2}, \quad \mathcal{K}(1) = \infty, \end{cases}$$

and

$$(1.2) \quad \begin{cases} \mathcal{E}(r) = \int_0^{\pi/2} \sqrt{1 - r^2 \sin^2 \theta} d\theta, \\ \mathcal{E}'(r) = \mathcal{E}(r'), \\ \mathcal{E}(0) = \frac{\pi}{2}, \quad \mathcal{E}(1) = 1. \end{cases}$$

In the sequel, we will use the symbols \mathcal{K} and \mathcal{E} for $\mathcal{K}(r)$ and $\mathcal{E}(r)$, respectively. The complete elliptic integrals play a very important role in the study of conformal invariants [7], quasiconformal mappings [5, 6, 7, 12] and Ramanujan's modular equations [4]. Numerous sharp inequalities and elementary approximations for the complete elliptic integrals have been proved in [2, 3, 7, 8, 13, 14].

The special function $m(r)$ is defined as

$$m(r) = \frac{2}{\pi} r'^2 \mathcal{K}(r) \mathcal{K}'(r), \quad 0 < r < 1.$$

This function is of importance in the research of distortion theory of quasiconformal mappings in the plane. Recently, various interesting inequalities of $m(r)$ have been obtained by several authors, see [3, 5, 6, 7, 15].

The power mean is defined for $x, y > 0$ and real parameter λ by

$$M_\lambda(x, y) = \left(\frac{x^\lambda + y^\lambda}{2} \right)^{1/\lambda} \quad \text{for } \lambda \neq 0, \text{ and } M_0(x, y) = \sqrt{xy}.$$

It is well known that $M_\lambda(x, y)$ is continuous and increasing with respect to λ . Many interesting properties of power means are given in [10]. Power mean inequalities for some special functions can be found in [1, 7, 16, 17, 18].

In this paper, we shall show a power mean inequality for the special function $m(r)$. Our main result is the following theorem:

1.3. Theorem. *Let λ be a real number. The inequality*

$$(1.4) \quad M_\lambda(m(x), m(y)) \leq m(M_\lambda(x, y))$$

holds for all $x, y \in (0, 1)$ if and only if $\lambda \leq 0$. The reverse of (1.4) holds for all $x, y \in (0, 1)$ if and only if $\lambda \geq C > 0$, where C is some constant. The sign of equality is valid in (1.4) if and only if $x = y$.

2. LEMMAS

In order to prove our main result we need some lemmas, which we present in this section. We establish some properties of certain functions, which are defined in terms of the complete elliptic integrals of the first and second kinds, \mathcal{K} and \mathcal{E} , respectively.

Now we list some derivative formulas [7, Appendix E, pp. 474-475]:

$$\frac{d\mathcal{K}}{dr} = \frac{\mathcal{E} - r'^2\mathcal{K}}{rr'^2}, \quad \frac{d\mathcal{E}}{dr} = \frac{\mathcal{E} - \mathcal{K}}{r},$$

and

$$\frac{d}{dr}m(r) = \frac{\pi - 4\mathcal{E}'\mathcal{K}}{\pi r},$$

where $0 < r < 1$. By the derivative of $m(r)$, it is easy to see that $m(r)$ is strictly decreasing from $(0, 1)$ onto $(0, \infty)$.

The following Lemma 2.1 is from [4, Lemma 5.2(2)] and [7, Theorem 3.21 (1), (7), and Exercises 3.43 (32)].

- 2.1. Lemma.** (1) *The function $f_1(r) = r'^2\mathcal{K}/\mathcal{E}$ is decreasing from $(0, 1)$ onto $(0, 1)$.*
 (2) *The function $f_2(r) = (\mathcal{E} - r'^2\mathcal{K})/r^2$ is strictly increasing and convex from $(0, 1)$ onto $(\pi/4, 1)$.*
 (3) *For each $c \in [1/2, \infty)$, the function $f_3(r) = r'^c\mathcal{K}$ is decreasing from $[0, 1)$ onto $(0, \pi/2]$.*
 (4) *The function $f_4(r) = r^{-2}(\mathcal{K} - \mathcal{E})/\mathcal{K}$ is increasing and convex from $(0, 1)$ onto $(1/2, 1)$.*

2.2. Lemma. *For $0 < r < 1$, let*

$$g(r) = \frac{\mathcal{K}\mathcal{K}'(\mathcal{E}\mathcal{E}' + r^2\mathcal{K}\mathcal{K}' - \mathcal{K}\mathcal{E}')}{(4\mathcal{E}'\mathcal{K} - \pi)^2}.$$

Then $g(r) > 0$ for all $r \in (0, 1)$, and $g(0^+) = g(1^-) = 0$.

Proof. By the formula (1.2) and the parts (1) and (2) of Lemma 2.1, we have

$$\frac{(4\mathcal{E}'\mathcal{K} - \pi)^2}{\mathcal{K}\mathcal{K}'}g(r) = \mathcal{E} \left(\mathcal{E}' - \frac{r'^2\mathcal{K}}{\mathcal{E}} \frac{\mathcal{E}' - r^2\mathcal{K}'}{r'^2} \right) > 0,$$

and hence $g(r) > 0$ for all $r \in (0, 1)$.

By Lemma 2.1 (3) and (4), we get

$$\begin{aligned} g(0^+) &= \lim_{r \rightarrow 0^+} g(r) = \lim_{r \rightarrow 1^-} \frac{\mathcal{K}\mathcal{K}'(\mathcal{E}\mathcal{E}' + r'^2\mathcal{K}\mathcal{K}' - \mathcal{K}'\mathcal{E})}{(4\mathcal{E}\mathcal{K}' - \pi)^2} \\ &= \lim_{r \rightarrow 1^-} \frac{\mathcal{K}'}{(4\mathcal{E}\mathcal{K}' - \pi)^2} \left[(r'\mathcal{K})^2\mathcal{K}' - \frac{\mathcal{K}' - \mathcal{E}'}{r'^2\mathcal{K}'}(r'^2\mathcal{K})(\mathcal{K}'\mathcal{E}) \right] \\ &= 0, \end{aligned}$$

and

$$g(1^-) = \lim_{r \rightarrow 1^-} g(r) = \lim_{r \rightarrow 1^-} \frac{\mathcal{K}'}{(4\mathcal{E}' - \pi/\mathcal{K})^2} \left(\frac{\mathcal{E}\mathcal{E}'}{\mathcal{K}} + r^2\mathcal{K}' - \mathcal{E}' \right) = 0.$$

□

2.3. Lemma. *Let λ be a real number. The function*

$$h(r) = \frac{4\mathcal{E}'\mathcal{K} - \pi}{m(r)} \left(\frac{m(r)}{r} \right)^\lambda$$

is strictly increasing on $(0, 1)$ if and only if $\lambda \leq 0$.

Proof. By logarithmic differentiation,

$$\begin{aligned} \frac{h'(r)}{h(r)} &= \frac{4 \left[\mathcal{K} \frac{\mathcal{E}' - \mathcal{K}'}{r'} \left(-\frac{r}{r'} \right) + \mathcal{E}' \frac{\mathcal{E} - r'^2\mathcal{K}}{rr'^2} \right]}{4\mathcal{E}'\mathcal{K} - \pi} - \frac{\pi - 4\mathcal{E}'\mathcal{K}}{\pi r m(r)} + \lambda \left(\frac{\pi - 4\mathcal{E}'\mathcal{K}}{\pi r m(r)} - \frac{1}{r} \right) \\ &= \left(\frac{4\mathcal{E}'\mathcal{K} - \pi}{\pi r m(r)} + \frac{1}{r} \right) \times \\ (2.4) \quad &\left[\frac{1}{1 + \frac{\pi m(r)}{4\mathcal{E}'\mathcal{K} - \pi}} \left(1 + 8 \frac{\mathcal{K}\mathcal{K}'(\mathcal{E}\mathcal{E}' + r^2\mathcal{K}\mathcal{K}' - \mathcal{K}\mathcal{E}')}{(4\mathcal{E}'\mathcal{K} - \pi)^2} \right) - \lambda \right], \end{aligned}$$

which is positive for all $r \in (0, 1)$ if and only if $\lambda \leq 0$ by Lemma 2.2, since $m(r)/(4\mathcal{E}'\mathcal{K} - \pi)$ is clearly decreasing from $(0, 1)$ onto $(0, \infty)$. □

2.5. Remark. *Let*

$$H(r) = \frac{1}{1 + \frac{\pi m(r)}{4\mathcal{E}'\mathcal{K} - \pi}} \left(1 + 8 \frac{\mathcal{K}\mathcal{K}'(\mathcal{E}\mathcal{E}' + r^2\mathcal{K}\mathcal{K}' - \mathcal{K}\mathcal{E}')}{(4\mathcal{E}'\mathcal{K} - \pi)^2} \right)$$

for $r \in (0, 1)$ and $H(0) = 0$, $H(1) = 1$. Then H is continuous on $[0, 1]$. Hence there exists $r_0 \in (0, 1]$ such that H obtains its maximum C at r_0 . Thus it is easy to conclude from (2.4) that h is strictly decreasing on $(0, 1)$ if and only if $\lambda \geq C$.

2.6. Open problem. *What is the exact expression for C ?*

3. PROOF OF THE MAIN RESULT

We are now in a position to prove the main result.

Proof of Theorem 1.3. We first prove the inequality (1.4) for $\lambda \neq 0$. We may assume that $x \leq y$. Define

$$F(x, y) = m(M_\lambda(x, y))^\lambda - \frac{m(x)^\lambda + m(y)^\lambda}{2}, \quad \lambda \neq 0.$$

Let $t = M_\lambda(x, y)$, then $\frac{\partial t}{\partial x} = \frac{1}{2}(\frac{x}{t})^{\lambda-1}$. If $x < y$, we have that $t > x$. By differentiation,

$$\begin{aligned} \frac{\partial F}{\partial x} &= \frac{\lambda}{2} m(t)^{\lambda-1} \frac{\pi - 4\mathcal{E}'(t)\mathcal{K}(t)}{\pi t} \left(\frac{x}{t}\right)^{\lambda-1} - \frac{\lambda}{2} m(x)^{\lambda-1} \frac{\pi - 4\mathcal{E}'(x)\mathcal{K}(x)}{\pi x} \\ &= \frac{\lambda x^{\lambda-1}}{2\pi} \left[\frac{4\mathcal{E}'(x)\mathcal{K}(x) - \pi}{m(x)} \left(\frac{m(x)}{x}\right)^\lambda - \frac{4\mathcal{E}'(t)\mathcal{K}(t) - \pi}{m(t)} \left(\frac{m(t)}{t}\right)^\lambda \right] \end{aligned}$$

which is positive if and only if $\lambda < 0$ by Lemma 2.3. Hence $F(x, y)$ is strictly increasing with respect to x and $F(x, y) \leq F(y, y) = 0$. We now obtain the inequality

$$m(M_\lambda(x, y))^\lambda \leq \frac{m(x)^\lambda + m(y)^\lambda}{2},$$

that is $m(M_\lambda(x, y)) \geq M_\lambda(m(x), m(y))$ if and only if $\lambda < 0$, with the equality if and only if $x = y$.

Similarly, by the statement in Remark 2.5 one can see that the reverse of (1.4) holds for all $x, y \in (0, 1)$ if and only if $\lambda \geq C > 0$, C is the same as Remark 2.5, with the equality if and only if $x = y$.

Now we prove the inequality (1.4) for $\lambda = 0$. We may assume that $x \leq y$. Define

$$G(x, y) = \frac{m(\sqrt{xy})^2}{m(x)m(y)}.$$

Let $t = \sqrt{xy}$, then $\frac{\partial t}{\partial x} = \frac{1}{2} \frac{t}{x}$. If $x < y$, we have that $t > x$. By logarithmic differentiation, we have

$$\begin{aligned} \frac{1}{G(x, y)} \frac{\partial G}{\partial x} &= \frac{1}{m(t)} \frac{\pi - 4\mathcal{E}'(t)\mathcal{K}(t)}{\pi t} \frac{t}{x} - \frac{1}{m(x)} \frac{\pi - 4\mathcal{E}'(x)\mathcal{K}(x)}{\pi x} \\ &= \frac{1}{\pi x} \left(\frac{4\mathcal{E}'(x)\mathcal{K}(x) - \pi}{m(x)} - \frac{4\mathcal{E}'(t)\mathcal{K}(t) - \pi}{m(t)} \right) \end{aligned}$$

which is negative. Hence $G(x, y)$ is strictly decreasing with respect to x and $G(x, y) \geq G(y, y) = 1$. We now obtain the inequality

$$m(\sqrt{xy}) \geq \sqrt{m(x)m(y)},$$

that is $m(M_0(x, y)) \geq M_0(m(x), m(y))$, with the equality if and only if $x = y$. This completes the proof. \square

The following corollary is clear.

3.1. Corollary. *The function $m(r)$ is concave on $(0, 1)$ with respect to power mean of order λ if and only if $\lambda \leq 0$, and convex on $(0, 1)$ with respect to power mean of order λ if and only if $\lambda \geq C$.*

Acknowledgments. This research is partly supported by the NNSF of China (No. 11071069) and the NSF of Zhejiang Province (No. Y6100170).

REFERENCES

- [1] H. Alzer, *A power mean inequality for the gamma function*, Monatsh. Math. **131**(2000), 179-188.
- [2] H. Alzer and S.-L. Qiu, *Monotonicity theorems and inequalities for the complete elliptic integrals*, J. Comput. Appl. Math. **172**(2004), 289-312.
- [3] G.D. Anderson, S.-L. Qiu and M.K. Vamanamurthy, *Elliptic integral inequalities, with applications*, Constr. Approx. **14**(1998), 195-207.
- [4] G.D. Anderson, S.-L. Qiu, M.K. Vamanamurthy and M. Vuorinen, *Generalized elliptic integrals and modular equations*, Pac. J. Math. **192**(2000), 1-37.
- [5] G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen, *Distortion functions for plane quasiconformal mappings*, Israel J. Math. **62**(1988), 1-16.
- [6] G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen, *Inequalities for plane quasiconformal mappings*, Contemp. Math. 169, Amer. Math. Soc., 1994, 1-27.
- [7] G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [8] R.W. Barnard, K. Pearce and K.C. Richards, *An inequality involving the generalized hypergeometric function and the arc length of an ellipse*, SIAM J. Math. Anal. **31**(2000), 693-699.
- [9] F. Bowman, *Introduction to Elliptic Functions with Applications*, Dover, New York, 1961.
- [10] P.S. Bullen, *Handbook of Means and Their Inequalities*, Mathematics and Its Applications 560, Kluwer Academic Publishers, Dordrecht, 2003.
- [11] P.F. Byrd and M.D. Friedman, *Handbook of Elliptic integrals for Engineers and Physicists*, 2nd ed., Die Grundlehren Math. Wiss. 67, Springer-Verlag, Berlin, 1971.
- [12] O. Hübner, *Remarks on a paper of Lawrynowicz on quasiconformal mappings*, Bull. de L'Acad. Polon. des Sci. **18**(1970), 183-186.
- [13] H. Kazi and E. Neuman, *Inequalities and bounds for elliptic integrals. II*, Special functions and orthogonal polynomials, 127-138, Contemp. Math. 471, Amer. Math. Soc., Providence, RI, 2008.
- [14] H. Kazi and E. Neuman, *Inequalities and bounds for elliptic integrals*, J. Approx. Theory **146**(2007), 212-226.
- [15] S.-L. Qiu and M. Vuorinen, *Infinite products and normalized quotients of hypergeometric functions*, SIAM J. Math. Anal. **30**(1999), 1057-1075.
- [16] K.C. Richards, *Sharp power mean bounds for the Gaussian hypergeometric function*, J. Math. Anal. Appl. **308**(2005), 303-313.
- [17] G.D. Wang, X.H. Zhang and Y.P. Jiang, *Concavity with respect to Hölder means involving the generalized Grötzsch function*, J. Math. Anal. Appl. **379**(2011), 200-204.
- [18] X.H. Zhang, G.D. Wang and Y.M. Chu, *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl. **353**(2009), 256-259.

COLLEGE OF MATHEMATICS AND ECONOMETRICS, HUNAN UNIVERSITY, CHANGSHA 410082, CHINA

E-mail address: wgdwang@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TURKU, FI-20014 TURKU, FINLAND

E-mail address: xiazha@utu.fi

DEPARTMENT OF MATHEMATICS, HUZHOU TEACHERS COLLEGE, HUZHOU 313000, CHINA

E-mail address: chuyuming@hutc.zj.cn